

evidentia.ch – Wissen und Werkzeuge für Ärzte

# Online Marketing von Medikamenten

Zürich, 12. Januar 2017

## Intro






## Ärzte online – ein paar Fakten



## Lösungsansätze



→ Herausforderung: Kreation von geeigneten und für den Arzt/Ärztin nützlichen Inhalten für den Online-Kanal

## Case „Slide Kit“

ESSENTIAL SLIDE KIT

# DIAGNOSE UND THERAPIE DER IDIOPATHISCHEN LUNGENFIBROSE (IPF)



Autor: PD Dr. med. Christian Clarenbach, Zürich  
Autor: Dr. med. Manuela Funke-Chambour, Bern

Letzte Aktualisierung: 28. April 2016

Mit freundlicher Unterstützung\* von:



\* Dieses Essential Slide Kit wurde von einem unabhängigen Team von Pneumologen erstellt und dient ausschließlich Informationszwecken.





evidentia  
Wissen und Werkzeuge für Ärzte

## Case „Rechner“

57	kg	172	cm
Berechneter BMI Score		19	


BMI (kg/m <sup>2</sup> )	Internationale Klassifikation nach WHO <sup>1</sup>
< 18.5	Untergewicht
18.5 - 25	Normalgewicht
25 - 30	Übergewicht (Präadipositas)
30 - 35	Adipositas Grad I
35 - 40	Adipositas Grad II
≥ 40	Adipositas Grad III

## Case „evidentia Kompendium“

Präparat	Wirkstoffe	Galenische Form	Stärke	Packungsgrösse	EFP	PP/- Vergleich	SB	Zulassungs- inhaber	Kategori- sierung
Levetiracetam (N03AX14) 86 Präparate									
<a href="#">Kepra 250 mg</a>	Levetiracetamum	Filmtabletten	250 mg	30 Tablette(n)	14.99	<u>29.50</u>	10%	UCB-Pharma SA	B / SL / SO
<a href="#">Kepra 500 mg</a>	Levetiracetamum	Filmtabletten	500 mg	20 Tablette(n)	15.26	<u>33.90</u>	10%	UCB-Pharma SA	B / SL / SO
<a href="#">Kepra 500 mg</a>	Levetiracetamum	Filmtabletten	500 mg	100 Tablette(n)	79.10	<u>107.20</u>	10%	UCB-Pharma SA	B / SL / SO
<a href="#">Kepra 500 mg</a>	Levetiracetamum	Filmtabletten	500 mg	2 x 100 Tablette(n)	154.18	<u>193.40</u>	10%	UCB-Pharma SA	B / SL / SO
<a href="#">Kepra 1000 mg</a>	Levetiracetamum	Filmtabletten	1000 mg	30 Tablette(n)	49.88	<u>73.65</u>	10%	UCB-Pharma SA	B / SL / SO
<a href="#">Kepra 1000 mg</a>	Levetiracetamum	Filmtabletten	1000 mg	100 Tablette(n)	152.87	<u>191.90</u>	10%	UCB-Pharma SA	B / SL / SO
<a href="#">Kepra 1000 mg</a>	Levetiracetamum	Filmtabletten	1000 mg	2 x 100 Tablette(n)	295.43	<u>355.55</u>	10%	UCB-Pharma SA	B / SL / SO
<a href="#">Kepra 100 mg/mL</a>	Levetiracetamum	Lösung (oral)	100 mg/ml	Flasche(n) à 300 ml	56.94	<u>81.75</u>	10%	UCB-Pharma SA	B / SL / SO
<a href="#">Levetiracetam Desitin 250 mg</a>	 Levetiracetamum	Filmtabletten	250 mg	30 Tablette(n)	11.87	<u>25.95</u>	10%	Desitin Pharma GmbH	B / SL / SG
<a href="#">Levetiracetam Desitin 500 mg</a>	 Levetiracetamum	Filmtabletten	500 mg	20 Tablette(n)	13.83	<u>28.20</u>	10%	Desitin Pharma GmbH	B / SL / SG
<a href="#">Levetiracetam Desitin 500 mg</a>	 Levetiracetamum	Filmtabletten	500 mg	100 Tablette(n)	63.53	<u>89.35</u>	10%	Desitin Pharma GmbH	B / SL / SG
<a href="#">Levetiracetam Desitin 500 mg</a>	 Levetiracetamum	Filmtabletten	500 mg	200 Tablette(n)	125.36	<u>160.30</u>	10%	Desitin Pharma GmbH	B / SL / SG



# Case „Product Channel“


×

Klinische Wirksamkeit
Dosierung und Sicherheit
Alles um Esbriet®
Ihr Kontakt

**Dosierungsempfehlung**

	Morning Breakfast	Afternoon Lunch	Evening Dinner
Days 1–7	x 1	x 1	x 1
Days 8–14	x 2	x 2	x 2
Days 15+	x 3	x 3	x 3

**Monitoring**

**Elevated liver enzymes\***

**Monitoring**

- ALT, AST, and bilirubin levels should be measured prior to the initiation of therapy, monthly for the first 6 months, and every 3 months thereafter
- Patients should be instructed to report symptoms of liver disease (eg, dark urine or jaundice) to their physician


Enzyme level	Dose modification
ALT and/or AST >3x but <5x ULN without symptoms or hyperbilirubinemia	<ul style="list-style-type: none"> <li>• Discontinue any confounding medications, exclude other causes, and monitor patient closely</li> <li>• Repeat liver function tests as clinically indicated</li> <li>• The full dosage of Esbriet may be maintained, if clinically appropriate, or reduced or interrupted</li> <li>• Dosing can be retitrated to full dosage as tolerated</li> </ul>
ALT and/or AST >3x but <5x ULN with symptoms or hyperbilirubinemia or ALT and/or AST >5x ULN regardless of symptoms or hyperbilirubinemia	<ul style="list-style-type: none"> <li>• Permanently discontinue Esbriet and do not rechallenge</li> </ul>

ALT—alanine aminotransferase; AST—aspartate aminotransferase; ULN—upper limit of normal.


**Nebenwirkungen**

ASCEND, CAPACITY 004, and CAPACITY 006 trials: Adverse reactions occurring in ≥10% of Esbriet-treated patients and more commonly than with placebo	% of patients (0–118 weeks)	
	Esbriet (n=620)	Placebo (n=624)
Nausea	36	16
Rash	30	10
Abdominal pain*	24	15
Upper respiratory tract infection	27	25
Diarrhea	28	20
Fatigue	26	19
Headache	22	19
Dyspepsia	19	7
Dizziness	18	11
Vomiting	13	6
Anorexia	13	5
Gastroesophageal reflux disease	11	7
Sinusitis	11	10
Insomnia	10	7
Weight decreased	10	5
Arthralgia	10	7
<b>Discontinuation due to adverse reactions</b>	<b>14.6</b>	<b>9.6</b>

\*includes abdominal pain, upper abdominal pain, abdominal distension, and stomach discomfort


×

Klinische Wirksamkeit
Dosierung und Sicherheit
Alles um Esbriet®
Ihr Kontakt



**Produktmanager Schweiz**

---

**Aussendienst Zentral- und Nordwestschweiz**

---

**Aussendienst Region Zürich und Ostschweiz**

Lena Muster  
+ 41 800 00 00  
lena.muster@entresto.ch

---

Sandro Muster  
+ 41 800 00 03  
sandro.muster@entresto.ch

---

Max Muster  
+ 41 800 00 04  
max.muster@entresto.ch

## Ihr Ansprechpartner

### **evidentia ag**

Seefeldstrasse 90

CH - 8008 Zürich

+41 44 380 54 46 (fon)

+41 44 380 54 45 (fax)

[www.evidentia.ch](http://www.evidentia.ch)

[info@evidentia.ch](mailto:info@evidentia.ch)

### **Beat Steinegger**

lic. oec. HSG, Managing Partner

Email: [beat.steinegger@evidentia.ch](mailto:beat.steinegger@evidentia.ch)

Mobile: +41 79 570 07 03